

- ✓ Foliar
- ✓ Liquid Injection
- ✓ Fertigation



Technical Brochure

Updated on: - 25 July 2011

Hi-2 Trace

Concentrate complex blend with copper & zinc chelated. Suitable for foliar, furrow injection and fertigation

Some of the benefits include:-

- Efficiently corrects plant deficiencies where Cu, Zn, S are lacking
- Easily combines with other nutrients
- Improved nutrition uptake due to stimulated root activity
- Improves resistance to disease as a result of increased brix levels
- Essential for grain fill

TYPICAL ANALYSIS	G/L	% W/V
Zinc Sulphate (Zn)	100	10
Copper Sulphate (Cu)	20	2.0
Sulphur (S)	59	5.9

Note: Due to sulphur in product, do a jam jar test if other products are added for liquid injecting systems.

APPLICATION

Foliar Application	Rate/Ha	Water
Broadacre	0.5-2L	50-100L
Horticulture	0.5-4L	300-1500L
Orchards (stone fruit excluded)	0.5-1L	1000L
Viticulture	0.5-3L	300-1200L
Furrow Injection	Rate/Ha	Water
Broadacre	2-5L	80L
Fertigation	Rate/Ha	Water
Horticulture, Viticulture, Orchards	5-10L	Variable

pH 4.05
SG 1.285

Important: - Do not foliar spray stone-fruit. Do not foliar spray any crop at the start of and during flowering

PACKAGING

10L, 20L, 200L and 1000L

24 Shanahan Road, Davenport, Bunbury WA, 6230 > PO Box 5351, Davenport, Bunbury, WA 6230 > www.hitechag.com.au

Ph (08) 9725 7322 Fax (08) 9725 7342 email: - sales@hitechag.com.au